



## IMAGE IN CARDIOLOGY

# Functional assessment of coronary disease using workstation-based computed tomography-derived fractional flow reserve



## Avaliação funcional de doença coronária utilizando FFR derivado de TC

Nuno Dias Ferreira\*, Ricardo Ladeiras-Lopes, Gustavo Pires de Moraes, Vasco Gama, Pedro Braga

*Serviço de Cardiologia, Centro Hospitalar de Vila Nova de Gaia/Espinho, Vila Nova de Gaia, Portugal*

Received 12 August 2018; accepted 10 September 2018

Available online 22 November 2019


A 55-year-old man presented to our outpatient clinic reporting a six-month history of exertional dyspnea and intermittent atypical chest pain. His medical history included essential hypertension. The baseline electrocardiogram and echocardiogram were normal. The patient underwent treadmill exercise stress testing under the Bruce protocol. During stage 3, the patient developed 1.5-mm horizontal ST depression in V3-V6 and the inferior leads. The test was interrupted at 9 min due to fatigue, with no chest pain.

Due to intermediate risk findings on the treadmill test, coronary computed tomography angiography (CCTA) was requested for further assessment. The coronary calcium score was 105.2 Agatston units and an intermediate stenosis was identified in the proximal left anterior descending artery (LAD) due to non-calcified plaque (Figure 1A and D,


arrow). Because the physiologic significance of the lesion was unclear, computed tomography-derived fractional flow reserve (CT-FFR) was determined using a workstation-based software prototype (cFFR, Siemens AG Healthcare, Forchheim, Germany). The value of CT-FFR just beyond the proximal LAD stenosis was 0.75 (Figure 2). The patient was referred for invasive coronary angiography (ICA) which confirmed a 75% stenosis in the proximal LAD (Figure 3B and Supplementary Video S1). Invasive FFR assessed in the LAD showed a similar result (0.76) to that observed on CCTA (Figures 3A and 3C). The patient successfully underwent percutaneous coronary intervention with a drug-eluting stent (Figure 4) and was asymptomatic after the procedure. This case illustrates the value of CT-FFR as a gatekeeper prior to ICA, to appropriately select patients and vessels for revascularization.

\* Corresponding author.


E-mail address: [ferreira.nmd@gmail.com](mailto:ferreira.nmd@gmail.com) (N. Dias Ferreira).




**Figure 1** Coronary computed tomography angiography images. Curved multiplanar reformats of the left anterior descending artery (LAD) (A), left circumflex artery (B) and right coronary artery (C); three-dimensional volume rendering image of the heart (D). Note the presence of significant stenosis in the proximal LAD due to non-calcified plaque (arrows).



**Figure 2** Computed tomography (CT)-derived fractional flow reserve (FFR) imaging showing an intermediate stenosis in the proximal left anterior descending artery. The computed value of CT-derived FFR was 0.75.



**Figure 3** Correlation between invasive coronary angiography and coronary computed tomography (CT) angiography. (A) CT-derived fractional flow reserve (FFR) imaging showing an intermediate stenosis in the proximal left anterior descending artery (LAD) with a computed FFR of 0.75; (B) invasive coronary angiography confirming a 75% stenosis in the proximal LAD; (C) the measured value of invasive FFR was 0.76, which showed a good agreement with the CT-derived value (A).



**Figure 4** Images from invasive coronary angiography before (A) and after (B) percutaneous coronary intervention in the proximal left anterior descending artery, with successful implantation of a drug-eluting stent.

### Conflicts of interest

The authors have no conflicts of interest to declare.

### Appendix A. Supplementary material

Supplementary material associated with this article can be found in the online version at [doi:10.1016/j.repc.2018.09.015](https://doi.org/10.1016/j.repc.2018.09.015).