



## EDITORIAL COMMENT

# Keep it as simple as possible<sup>☆</sup>

## Deixá-lo tão simples quanto possível

Rui Ferreira



*Serviço de Cardiologia, Hospital de Santa Marta, Centro Hospitalar de Lisboa Central, Lisboa, Portugal*

Available online 28 February 2018

Primary angioplasty as revascularization therapy to treat myocardial infarction (MI) significantly reduces mortality in patients with this condition, for which it is currently the treatment of choice.<sup>1,2</sup> MI is a true cardiovascular emergency, given the linear relationship between time since onset and outcome.

The decision to adopt this therapeutic strategy obviously requires the operator to have specific skills, which are strongly influenced by the annual volume of procedures performed at the center and by each operator individually. On the basis of multiple studies, recommended minimum volumes of procedures have been defined, which are now stipulated in the international guidelines.

From a technical standpoint, advances in technology and engineering have increased the availability of new materials with characteristics that enable procedures to be performed rapidly and effectively, including catheters with novel geometry, more maneuverable guidewires, new means of mechanical thrombus aspiration, and stents with better profiles.

In recent years, another important change has taken place: the use of radial access in preference to the previously standard femoral access. Various studies have shown radial access to be superior, mainly due to a significant reduction in bleeding complications.<sup>3,4</sup>

Several factors have thus come together to help achieve the main purpose of primary angioplasty, which is recanalization of the culprit artery and the fastest possible return to adequate flow (TIMI 3).

The accumulated experience of operators and teams have also enabled the use of certain techniques to shorten procedure time, such as the first-line use of a guidewire to approach the likely culprit artery (the right coronary artery in inferior MI or the left coronary in anterior MI).

The study by MC Arokiaraj published in this issue of the *Journal*<sup>5</sup> describes the experience in their center of a series of 34 cases with a different solution: the use of diagnostic catheters, with a larger internal diameter, for angioplasty, enabling the deployment of balloons and stents, as well as of catheters for intravascular ultrasound and optical coherence tomography.

This solution is recommended for emergency situations, particularly MI and acute coronary syndromes, in which selective catheterization of the culprit artery with appropriate guide catheters is more difficult, thus avoiding prolonged procedure time and the use of greater quantities of contrast agents and hence risk of nephrotoxicity.

As the author points out, this variation from the usual technique has significant practical limitations, notably when more complex maneuvers are foreseen, such as for treating bifurcations or when mechanical thrombus aspiration will be required.

It is also important to note that, in the patient group described here, radial access, in which the criteria for choice of catheters are different, was not used, and all the catheters were from a single manufacturer. It is

<sup>☆</sup> Please cite this article as: Ferreira R. Deixá-lo tão simples quanto possível. Rev Port Cardiol. 2018;37:127–128.

E-mail address: [cruzferreira@netcabo.pt](mailto:cruzferreira@netcabo.pt)

unclear whether the considerations presented can be generalized to the different equipment available at other laboratories.

Irrespective of the advantages that this solution may have in specific situations, we would highlight the crucial importance of the level of experience of centers and of operators performing primary angioplasty for MI and other acute coronary syndromes. This experience is directly related to the volume of procedures<sup>6</sup> and is without doubt an essential factor in obtaining good clinical outcomes.

### Conflicts of interest

The author has no conflicts of interest to declare.

### References

1. Ibanez B, James S, Agewall S, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST segment elevation: the Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2017, <http://dx.doi.org/10.1093/eurheartj/ehx393>.
2. O’Gara PT, Kushner FG, Ascheim DD, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61:e78–140.
3. Bernat I, Horak D, Stasek J, et al. ST-segment elevation myocardial infarction treated by radial or femoral approach in a multicenter randomized clinical trial: the STEMI-RADIAL trial. *J Am Coll Cardiol*. 2014;63:964–72 [Epub ahead of print].
4. Hinohara TT, Rao SV. Current state of radial artery catheterization in ST-elevation myocardial infarction. *Prog Cardiovasc Dis*. 2015;58:241–6 [review; Epub ahead of print].
5. Arokiaraj MC. Emergency coronary angioplasty with stenting using Cordis diagnostic coronary catheters when there is difficulty in engaging guide catheters and bench evaluation of diagnostic and guide catheters. *Rev Port Cardiol*. 2018;37:117–25.
6. Gutierrez A, Tsai TT, Stanislawski MA, et al. Adoption of transradial percutaneous coronary intervention and outcomes according to center radial volume in the Veterans Affairs Healthcare system: insights from the Veterans Affairs clinical assessment, reporting, and tracking (CART) program. *Circ Cardiovasc Interv*. 2013;6:336–46 [Epub ahead of print].