

IMAGES IN CARDIOLOGY

Left ventricular apical hypoplasia: An unusual diagnosis

Hipoplasia apical do ventrículo esquerdo: um diagnóstico raro

Alexandra Sousa^{a,*}, Teresa Pinho^a, Pedro Almeida^a, António Madureira^b,
Filipe Macedo^a, Maria Júlia Maciel^a

^a Cardiology Department, Centro Hospitalar de São João, Porto, Portugal

^b Radiology Department, Centro Hospitalar de São João, Porto, Portugal

Received 25 July 2012; accepted 6 August 2012

Available online 27 February 2013

A 36-year-old woman was referred for medical evaluation prior to non-cardiac surgery. She was healthy and only complained of sporadic palpitations. The 12-lead electrocardiogram showed sinus rhythm and left bundle branch block. Transthoracic echocardiography revealed a normally

functioning, slightly dilated left ventricle (LV), with the interventricular septum bulging toward the right ventricle (RV), which was elongated and wrapped around the LV (Figure 1; Supplementary material Videos S1 and S2). The papillary muscles were structurally abnormal and presented




Figure 1 Transthoracic echocardiogram. (A) Apical 4-chamber view; (B) apical 2-chamber view. Solid arrow: right ventricle. Dashed arrow: network of papillary muscles.

* Corresponding author.

E-mail address: xanasousa81@gmail.com (A. Sousa).




Figure 2 (A) Transthoracic echocardiogram, parasternal short-axis view. Contrast transthoracic echocardiogram: (B) parasternal short-axis view; (C) apical 4-chamber view; (D) apical 3-chamber view.

an apical origin, findings better visualized in contrast images (Figure 2; Videos S3–S5). Cardiac magnetic resonance confirmed the presence of a truncated LV, a complex network of papillary muscles of apical origin, and a banana-shaped, normally functioning RV (Figures 3 and 4; Videos S6–S8). There was no evidence of fatty infiltration or late enhancement with gadolinium.

During a three-year follow-up, the patient remained asymptomatic with no evidence of arrhythmias in serial Holter monitoring and with no signs of heart failure.

These findings are consistent with LV apical hypoplasia, a recently recognized cardiomyopathy, with only 15 cases reported since its first description in 2004. The age of diagnosis (3 months to 63 years) and clinical presentation vary




Figure 3 Cardiac magnetic resonance. Steady-state free precession images in horizontal long-axis view (A) and vertical long-axis view (B). Solid arrow: right ventricle. Dashed arrow: network of papillary muscles.



Figure 4 Cardiac magnetic resonance. Sequence of steady-state free precession images, in short-axis view, from base to mid-ventricular level (A–D), showing the complex network of papillary muscles.

widely, with most individuals remaining asymptomatic or mildly symptomatic for variable periods of time. However, a case of a 19-year-old male with a rapidly fatal presentation has been described. The precise pathophysiological basis and natural history of this entity remain unknown.

To our knowledge, this is the first case described in the Portuguese population.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the

publication of patient data and that all the patients included in the study received sufficient information and gave their written informed consent to participate in the study.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflicts of interest

The authors have no conflicts of interest to declare.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.repc.2012.08.011](https://doi.org/10.1016/j.repc.2012.08.011).