



IMAGE IN CARDIOLOGY

Patent foramen ovale may not always be benign in the elderly

Foramen oval patente pode não ser sempre benigno no idoso

Ana Aires^{a,b,*}, Nuno Oliveira^c, Filipe Macedo^{b,c}, Elsa Azevedo^{a,b}

^a Department of Neurology, São João Hospital Center, Porto, Portugal

^b Faculty of Medicine, University of Porto, Porto, Portugal

^c Department of Cardiology, São João Hospital Center, Porto, Portugal

Received 18 April 2018; accepted 14 May 2018

A 65-year-old woman was admitted to the hospital with fever and decompensated heart failure. She had long-lasting peripheral venous insufficiency and had been experiencing

heart failure symptoms for six months. She had no history of hypertension, diabetes mellitus, coronary heart disease, atrial fibrillation, stroke or transient ischemic attack.




Figure 1 A – Transesophageal echocardiography (TEE) reveals the passage of a large thrombus through the patent foramen ovale, which later led to stroke. B – TEE shows a smaller thrombus following the initiation of unfractionated heparin. C – Transthoracic echocardiography reveals no thrombus.

* Corresponding author.

E-mail address: ana.aires.mail@gmail.com (A. Aires).

Six days after admission, she developed acute pulmonary edema. A transesophageal echocardiography (TEE) revealed severe left ventricular dysfunction and an intracardiac thrombus measuring 5-6 cm, traversing a small patent foramen ovale (PFO) (Figure 1A). Unfractionated heparin was promptly initiated. Nevertheless, one week later, she woke up aphasic with right-sided hemiparesis. A brain computed tomography scan showed a left hemispheric stroke, while the patient's blood work, a cervical duplex scan and cardiac catheterization revealed no significant findings. The TEE was repeated and a smaller thrombus found (Figure 1B). Two months after discharge, the thrombus was undetectable (Figure 1C).

PFO is not considered to be a major cause of cardioembolic stroke, especially in the elderly, since other potential causes exist. Our case, however, shows that a causative

thrombus resulting from a paradoxical PFO embolism could be detected before giving rise to a stroke. This case is important as it allowed direct thrombus imaging within the PFO before it led to a stroke, which is not always possible, as well as revealing a smaller thrombus trapped inside the PFO after the stroke. The first thrombus visualized probably arose from the venous system of the lower limbs and passed through a small ostium, giving us the exceptional opportunity to prove the stroke's cardioembolic etiology through a PFO in an elderly patient.

Conflicts of interest

The authors have no conflicts of interest to declare.