

IMAGE IN CARDIOLOGY

A narrow path

Um trajeto estreito

Rodrigo Silva^{a,*}, Sofia Nogueira-Fernandes^a, Carina Arantes^a, João Maciel^b,
 Vítor Hugo Pereira^{a,c}, Catarina Vieira^a, Sérgia Rocha^a

^a Serviço de Cardiologia do Hospital de Braga, Braga, Portugal

^b Serviço de Cirurgia Cardiorádica do Centro Hospitalar de São João, Porto, Portugal

^c Escola de Medicina da Universidade do Minho, Braga, Portugal

Received 27 January 2025; accepted 24 March 2025

Available online 30 July 2025

A 35-year-old man was sent to the Cardiology consult due to recurrent syncope after orthostatism preceded by prodromal symptoms. His syncopal response to orthostatism was so significant and so fast that he resorted to wheelchair ambulation. Additionally, the patient also reported episodes of syncope while lying down. Physical examination revealed the presence of an exuberant pectus excavatum. He was admitted to the Cardiology Department for evaluation. The initial cardiac study did not reveal significant findings: the 12-lead electrocardiogram presented an incomplete right bundle branch block, electrocardiographic monitoring showed rare supraventricular ectopy, and the transthoracic echocardiogram presented no relevant findings. A computed tomography scan was performed which showed severe pectus excavatum (Figure 1), with the lower sternum distancing 6 mm from the vertebral column, compressing the inferior vena cava (arrow). This resulted in a Haller index (a pectus index obtained from dividing the transverse diameter of the chest by the anterior–posterior




Figure 1 A computed tomography scan axial slice revealing a severe case of pectus excavatum with an antero-posterior diameter of 6 mm and inferior vena cava compression (arrow).

* Corresponding author.

E-mail address: rjpintosilva@gmail.com (R. Silva).




Figure 2 A computed tomography scan axial slice after the Nuss procedure showing the post-operative outcome with an antero-posterior diameter of 49.5 mm.

diameter) of 46, far greater than the 3.25 cut-off to consider corrective surgery.¹ A cardiac magnetic resonance imaging was then performed, revealing severe inferior vena cava compression in decubitus position ([supplemental video](#)).

The patient was referred for thoracic surgery and underwent a Nuss procedure, minimally invasive surgery to correct this thoracic deformity using intrathoracic bars.² The patient did not experience any relevant complications from this, was syncope free during the follow-up and was able to walk once again. The chest radiograph and CT scan in follow-up revealed the excellent outcome of the surgery ([Figure 2](#)) with inferior vena cava decompression.

There are only a handful of published cases of pectus excavatum with inferior vena cava compression, with varying symptoms, such as syncopal episodes,³ lower extremity edema⁴ and pulsus paradoxus.⁵ To the best of our knowledge, we present the case with the most severe thoracic deformity causing disabling syncope in orthostatism, with a good functional outcome after surgical intervention.

Conflicts of interest

The authors have no conflicts of interest to declare.

Appendix A. Supplementary data

Supplementary material associated with this article can be found in the online version at <https://doi.org/10.1016/j.repc.2025.03.007>.

References

1. Haller JA Jr, Kramer SS, Lietman SA. Use of CT scans in selection of patients for pectus excavatum surgery: a preliminary report. *J Pediatr Surg.* 1987;22:904–6 [PMID: 3681619].
2. Velazco CS, Arsanjani R, Jaroszewski DE. Nuss procedure in the adult population for correction of pectus excavatum. *Semin Pediatr Surg.* 2018;27:161–9 [PMID: 30078487].
3. White JA, Fine NM, Shargall Y. Pectus excavatum with compression of the inferior vena cava: a rare cause of recurrent syncope. *Circulation.* 2009;120:1722–4.
4. Iannucci GJ, Slesnick TC, Kogon B, et al. Lower extremity edema in a child due to pectus excavatum. *Ann Thorac Surg.* 2015;99:e29–30 [PMID: 25639440].
5. Yalamanchili K, Summer W, Valentine V. Pectus excavatum with inspiratory inferior vena cava compression: a new presentation of pulsus paradoxus. *Am J Med Sci.* 2005;329:45–7 [PMID: 15654179].